

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number
WO 02/097039 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number: PCT/US02/16635

(22) International Filing Date: 23 May 2002 (23.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/294,076 29 May 2001 (29.05.2001) US

(71) Applicant: **LEXICON GENETICS INCORPORATED**
[US/US]; 4000 Research Forest Drive, The Woodlands, TX
77381 (US).

(72) Inventors: **YU, Xuanchuan**; 6 Danville Crossing, Conroe,
TX 77385 (US). **MIRANDA, Maricar**; 14814 Belterrazza
Drive, Houston, TX 77083 (US). **HU, Yi**; 333 Holly Creek
Ct. #203, The Woodlands, TX 77381 (US).

(74) Agents: **ISHIMOTO, Lance, K.** et al.; Lexicon Genetics
Incorporated, 4000 Research Forest Drive, The Woodlands,
TX 77381 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL HUMAN HYDROXYLASES AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in industrial, therapeutic, diagnostic, and pharmacogenomic applications.

WO 02/097039 A2

NOVEL HUMAN HYDROXYLASES AND
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S.
5 Provisional Application Number 60/294,076, which was
filed on May 29, 2001, and is herein incorporated by
reference in its entirety.

1. INTRODUCTION

10 The present invention relates to the discovery,
identification, and characterization of novel human
polynucleotides encoding proteins sharing sequence
similarity with mammalian hydroxylases. The invention
encompasses the described polynucleotides, host cell
15 expression systems, the encoded protein, fusion
proteins, polypeptides and peptides, antibodies to the
encoded proteins and peptides, and genetically
engineered animals that either lack or overexpress the
disclosed polynucleotides, antagonists and agonists of
20 the proteins, and other compounds that modulate the
expression or activity of the proteins encoded by the
disclosed polynucleotides, which can be used for
diagnosis, drug screening, clinical trial monitoring,
the treatment of diseases and disorders, and cosmetic or
25 nutraceutical applications.

2. BACKGROUND OF THE INVENTION

Hydroxylases are enzymes that mediate the cleavage
of hydroxyl groups, and many hydroxylases act in
30 biological pathways suitable for drug intervention.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery,
identification, and characterization of nucleotides that
35 encode novel human proteins, and the corresponding amino
acid sequences of these proteins. The novel human
proteins (NHPs) described for the first time herein

share structural similarity with animal hydroxylases, and particularly tryptophan hydroxylases, which are involved in a rate-limiting step in the biosynthesis of a number of neurologically active compounds, including,

5 but not limited to, DOPA, serotonin and melatonin.

Given the well known physiological functions of the above compounds, the described hydroxylases can be used to identify and/or develop agents useful for modulating behavior (i.e., treating anxiety, depression,

10 hyperactivity, sleep disorders, etc.). The novel human nucleic acid (cDNA) sequences described herein encode proteins/open reading frames (ORFs) of 490, 486, 485, and 484 amino acids in length (SEQ ID NOS:2, 4, 6, and 8, respectively).

15 In addition to the direct therapeutic use of the described NHPs, the invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHPs, peptides, and
20 antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and open reading frame or regulatory sequence replacement constructs) or to enhance the expression of the
25 described NHPs (e.g., expression constructs that place the described polynucleotide under the control of a strong promoter system), and transgenic animals that express a NHP sequence, or "knock-outs" (which can be conditional) that do not express a functional NHP.

30 Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cell ("ES cell") lines that contain gene trap mutations in a murine homolog of at least one of the described NHPs. When the unique NHP sequences described in SEQ ID NOS:1-
35 8 are "knocked-out" they provide a method of identifying phenotypic expression of the particular gene, as well as

a method of assigning function to previously unknown genes. In addition, animals in which the unique NHP sequences described in SEQ ID NOS:1-8 are "knocked-out" provide a unique source in which to elicit antibodies to homologous and orthologous proteins, which would have been previously viewed by the immune system as "self" and therefore would have failed to elicit significant antibody responses.

Additionally, the unique NHP sequences described in SEQ ID NOS:1-8 are useful for the identification of protein coding sequences, and mapping a unique gene to a particular chromosome. These sequences identify biologically verified exon splice junctions, as opposed to splice junctions that may have been bioinformatically predicted from genomic sequence alone. The sequences of the present invention are also useful as additional DNA markers for restriction fragment length polymorphism (RFLP) analysis, and in forensic biology, particularly given the presence of a nucleotide polymorphism within the described sequences.

Further, the present invention also relates to processes for identifying compounds that modulate, i.e., act as agonists or antagonists of, NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP products, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequences of the ORFs encoding the described NHP amino acid sequences.

5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs described for the first time herein are novel proteins that can be expressed in, *inter alia*, human cell lines, human fetal brain, spinal cord, lymph node, prostate, testis, thyroid, pancreas, pericardium, hypothalamus, fetal kidney, fetal lung, 6 and 9-week old embryos, osteosarcoma, and embryonic carcinoma cells.

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described polynucleotides, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of a NHP that correspond to functional domains of a NHP, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of a described NHP, in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, soluble proteins and peptides; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains (e.g., a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides, such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs, comprising a sequence first disclosed in the Sequence Listing.

As discussed above, the present invention includes the human DNA sequences presented in the

Sequence Listing (and vectors comprising the same), and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1x SSC/0.1% SDS at 68°C (Ausubel et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., N.Y., at p. 2.10.3) and encodes a functionally equivalent expression product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of a DNA sequence that encodes and expresses an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2x SSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species, and mutant NHPs, whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent Nos. 5,837,458 and 5,723,323 both of which are herein incorporated by reference in their entirety). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding a NHP ORF, or its functional equivalent, encoded by a polynucleotide sequence that is about 99, 95, 90, or about 85 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence

analysis package, as described herein, using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are
5 therefore the complements of, the described NHP nucleotide sequences. Such hybridization conditions may be highly stringent or less highly stringent, as described herein. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such
10 molecules are generally about 16 to about 100 bases long, or about 20 to about 80 bases long, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence
15 Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be
20 used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a microarray or high-throughput "chip" format). Additionally, a series of NHP oligonucleotide sequences, or the complements thereof, can be used to represent all
25 or a portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS:1-8 can be used as a hybridization probe in conjunction with a solid support
30 matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (i.e., gene chips, microtiter plates, etc.) of oligonucleotides and
35 polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers

present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed in at least one of the sequences of SEQ ID NOS:1-8, or an amino acid sequence encoded thereby.

5 Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon, are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186,
10 and 4,689,405, the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-8 can be used to identify and characterize the temporal and tissue specific expression
15 of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length of these probes is usually within a range of between about 8 to
20 about 2000 nucleotides. Preferably the probes consist of 60 nucleotides, and more preferably 25 nucleotides, from the sequences first disclosed in SEQ ID NOS:1-8.

For example, a series of such NHP oligonucleotide sequences, or the complements thereof, can be used in
25 chip format to represent all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length, can partially overlap each other, and/or the sequence may be
30 represented using oligonucleotides that do not overlap. Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in
35 the described Sequence Listing. Such oligonucleotide sequences can begin at any nucleotide present within a

sequence in the Sequence Listing, and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of
5 broad patterns of genetic activity, providing new understanding of gene functions, and generating novel and unexpected insight into transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ ID
10 NOS:1-8 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the identification of novel components, or gene functions that manifest themselves as novel phenotypes.

15 Probes consisting of sequences first disclosed in SEQ ID NOS:1-8 can also be used in the identification, selection, and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug targets, and
20 recognition of drug dependent changes in gene expression that are modulated through pathways distinct from the intended target of the drug. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

25 As an example of utility, the sequences first disclosed in SEQ ID NOS:1-8 can be utilized in microarrays, or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can
30 also be carried out using the sequences first disclosed in SEQ ID NOS:1-8 *in silico*, and by comparing previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-
35 8 can be used to identify mutations associated with a

particular disease, and also in diagnostic or prognostic assays.

Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be described by the net composition of the nucleotides present within a given region of the sequence, in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in SEQ ID NOS:1-8. Alternatively, a restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences, can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6x SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP antisense molecules, useful, for example, in NHP gene regulation and/or as antisense primers in amplification reactions of NHP nucleic acid sequences. With respect to NHP gene regulation, such techniques can be used to regulate biological functions.

Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded

- 5 oligonucleotides can additionally comprise at least one modified base moiety that is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine,
- 10 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,
- 15 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil,
- 20 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-
- 25 2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose,

- 30 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a

35 phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a

methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g., by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451), etc.

Low stringency conditions are well-known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (and periodic updates thereof), and Ausubel et al., 1989, *supra*.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The

identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

For example, the present sequences can be used in restriction fragment length polymorphism (RFLP) analysis to identify specific individuals. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification (as generally described in U.S. Patent No. 5,272,057, incorporated herein by reference). In addition, the sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e., another DNA sequence that is unique to a particular individual). Actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments.

Further, a NHP homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be genomic DNA, or total RNA, mRNA,

and/or cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known to express, or suspected of expressing, an allele of a NHP gene.

5 The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment
10 can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

 PCR technology can also be used to isolate full
15 length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (*i.e.*, one known to express, or suspected of expressing, a NHP gene, such as, for example, testis tissue). A reverse
20 transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase
25 reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see, *e.g.*, Sambrook
30 et al., 1989, *supra*.

 A cDNA encoding a mutant NHP sequence can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue
35 known to express, or suspected of expressing, a NHP, in an individual putatively carrying a mutant NHP allele,

and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal sequence.

5 Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well-known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a
10 corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of
15 carrying, or known to carry, a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, neurological disorders, such as, but not limited to, depression, anxiety, Alzheimer's disease or Parkinson's disease, obesity, high blood pressure,
20 connective tissue disorders, infertility, developmental disorders, etc.), or a cDNA library can be constructed using RNA from a tissue known to express, or suspected of expressing, a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled
25 and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well-known to those skilled in the art.

30 Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known to express, or suspected of expressing, a mutant NHP allele in an individual suspected of carrying, or known to carry,
35 such a mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed.

and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below (for screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor, N.Y.).

Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expression product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP expression product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well-known in the art.

The invention also encompasses: (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculovirus as described in U.S. Patent No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (i.e., gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators, and other

elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include, but are not limited to, the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 or adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP sequence (transcription factor inhibitors, antisense and ribozyme molecules, or open reading frame sequence or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs, or inappropriately expressed NHPs, for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of a NHP in the body. The use of engineered host cells and/or animals may

offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor for a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains corresponding to a NHP, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-mediated pathway), can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of a soluble NHP (or, for example, a suitably derivatized NHP, e.g., with polyethylene glycol (PEG), albumin, etc.), a NHP-IgFc fusion protein, or an anti-idiotypic antibody (or its Fab) that mimics the NHP, could activate or effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules, can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

5.1 THE NHP SEQUENCES

The cDNA sequences and the corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing. The described sequences were
5 compiled from human gene trapped sequences, and cDNAs made from human fetal brain, fetal lung, and lymph node mRNAs (Edge Biosystems, Gaithersburg, MD). As biologically validated (spliced and polyadenylated sequences), the described sequences are useful for
10 mapping the corresponding coding region of the human genome and particularly for mapping exon splice junctions. The described NHPs are apparently encoded on human chromosome 12 (see GENBANK accession no. AC023966).

15 An A/T polymorphism was detected during the sequencing of the NHPs at the nucleotide position represented by, for example, position 1125 of SEQ ID NO:1 (and the corresponding location in the other NHP nucleotide sequences), both of which result in an ala at
20 the region corresponding to amino acid (aa) position 375 of, for example, SEQ ID NO:2 (and the corresponding location in the other NHP amino acid sequences). As these polymorphisms are coding single nucleotide polymorphisms (SNPs), they are particularly useful in
25 forensic analysis.

An additional application of the described novel human polynucleotide sequences is their use in the molecular mutagenesis/evolution of proteins that are at least partially encoded by the described novel sequences
30 using, for example, polynucleotide shuffling or related methodologies. Such approaches are described in U.S. Patent Nos. 5,830,721 and 5,837,458, which are herein incorporated by reference in their entirety.

NHP gene products can also be expressed in
35 transgenic animals. Animals of any species, including, but not limited to, worms, mice, rats, rabbits, guinea

pigs, pigs, micro-pigs, birds, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees, may be used to generate NHP transgenic animals.

Any technique known in the art may be used to
5 introduce a NHP transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ
10 lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci. USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer
15 (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic
20 animals that carry a NHP transgene in all their cells, as well as animals that carry a transgene in some, but not all their cells, i.e., mosaic animals or somatic cell transgenic animals. A transgene may be integrated as a single transgene, or in concatamers, e.g., head-to-
25 head tandems or head-to-tail tandems. A transgene may also be selectively introduced into and activated in a particular cell-type by following, for example, the teaching of Lasko et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236. The regulatory sequences required for
30 such a cell-type specific activation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

When it is desired that a NHP transgene be integrated into the chromosomal site of the endogenous
35 NHP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing

some nucleotide sequences homologous to the endogenous NHP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous NHP gene (i.e., "knockout" animals).

The transgene can also be selectively introduced into a particular cell-type, thus inactivating the endogenous NHP gene in only that cell-type, by following, for example, the teaching of Gu et al., 1994, Science 265:103-106. The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant NHP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR. Samples of NHP gene-expressing tissue may also be evaluated immunocytochemically using antibodies specific for the NHP transgene product.

The present invention also provides for "knock-in" animals. Knock-in animals are those in which a polynucleotide sequence (i.e., a gene or a cDNA) that the animal does not naturally have in its genome is inserted in such a way that it is expressed. Examples include, but are not limited to, a human gene or cDNA used to replace its murine ortholog in the mouse, a murine cDNA used to replace the murine gene in the

mouse, and a human gene or cDNA or murine cDNA that is tagged with a reporter construct used to replace the murine ortholog or gene in the mouse. Such replacements can occur at the locus of the murine ortholog or gene, or at another specific site. Such knock-in animals are useful for the *in vivo* study, testing and validation of, *intra alia*, human drug targets, as well as for compounds that are directed at the same, and therapeutic proteins.

5.2 NHPS AND NHP POLYPEPTIDES

NHPS, NHP polypeptides, NHP peptide fragments, mutated, truncated, or deleted forms of the NHPS, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as therapeutic products (particularly in the treatment of behavioral disorders such as, but not limited to, dementia, insomnia, depression, anorexia, and premenstrual syndrome, etc., as well as cancers such as, but not limited to, lymphoma, leukemia, sarcomas, carcinomas, myelomas, etc.), as reagents in diagnostic assays, for the identification of other cellular gene products related to a NHP, as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and disease. Given the similarity information and expression data, the described NHPS can be targeted (by drugs, oligos, antibodies, etc.) in order to treat disease, or to therapeutically augment the efficacy of therapeutic or chemotherapeutic agents. In addition, the described NHPS can be used in drug screening assays similar to those described in, for example, U.S. Patent No. 6,048,850, herein incorporated by reference, in order to identify compounds for treating diseases such as, for

example, depression, anxiety, immune disorders, Alzheimer's disease, epilepsy, and Parkinson's disease.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP polynucleotides.

5 The NHPs display initiator methionines in DNA sequence contexts consistent with translation initiation sites, and a N-terminal signal sequence characteristic of secreted or membrane proteins (the signal sequence is often cleaved from the mature form of the protein
10 incident to the protein translocating across the membrane).

The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing, as well as analogues and derivatives
15 thereof, as well as any oligopeptide sequence of at least about 10-40 amino acids, about 12-35 amino acids, or about 16-30 amino acids in length first disclosed in the Sequence Listing. Further, corresponding NHP homologues from other species are encompassed by the
20 invention. In fact, any NHP encoded by the NHP nucleotide sequences described herein are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The
25 degenerate nature of the genetic code is well-known, and, accordingly, each amino acid presented in the Sequence Listing is generically representative of the well-known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such,
30 as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al., eds., Scientific American Books, New York, N.Y.,
35 herein incorporated by reference), are generically representative of all the various permutations and

combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences, as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described herein, but that result in a silent change, thus producing a functionally equivalent expression product. Amino acid substitutions can be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP products or NHP polypeptides can be produced in soluble or secreted forms (by removing one or more transmembrane domains where applicable), the peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells

that express a NHP, or a functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well-known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of a NHP, but to assess biological activity, e.g., in certain drug screening assays.

The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP nucleotide sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing NHP nucleotide sequences and promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is

to be produced for the generation of pharmaceutical compositions of or containing a NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in-frame with the *lacZ* coding region so that a fusion protein is produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Pharmacia or American Type Culture Collection) can also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target expression product can be released from the GST moiety.

In an exemplary insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign polynucleotide sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into a non-essential region (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of a NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted

sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an
5 adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the
10 adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., see Logan and Shenk,
15 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire
20 NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous
25 translational control signals, including, perhaps, the ATG initiation codon, may be provided. Furthermore, the initiation codon should be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous
30 translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bitter et
35 al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the expression product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and expression products. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for the desired processing of the primary transcript, glycosylation, and phosphorylation of the expression product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the NHP sequences described herein can be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci, which in turn can be cloned and expanded into cell lines. This method

may advantageously be used to engineer cell lines that express a NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of a NHP product.

A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes, which can be employed in tk⁻, hgp⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., 1980, Proc. Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hyg^r, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. Another exemplary system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the sequence of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are

loaded onto Ni^{2+} -nitriloacetic acid-agarose columns, and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are
5 fusion proteins that direct a NHP to a target organ and/or facilitate transport across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching an appropriate
10 signal sequence to a NHP would also transport a NHP to a desired location within the cell. Alternatively targeting of a NHP or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in "Liposomes:
15 A Practical Approach", New, R.R.C., ed., Oxford University Press, N.Y., and in U.S. Patent Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective disclosures, which are herein incorporated by reference in their entirety. Additionally embodied are
20 novel protein constructs engineered in such a way that they facilitate transport of NHPs to a target site or desired organ, where they cross the cell membrane and/or the nucleus where the NHPs can exert their functional activity. This goal may be achieved by coupling of a
25 NHP to a cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S. Provisional Patent Application Ser. Nos. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples
30 of such transducing sequences), to facilitate passage across cellular membranes, and can optionally be engineered to include nuclear localization signals.

Additionally contemplated are oligopeptides that are modeled on an amino acid sequence first described in
35 the Sequence Listing. Such NHP oligopeptides are generally between about 10 to about 100 amino acids

long, or between about 16 to about 80 amino acids long, or between about 20 to about 35 amino acids long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such NHP oligopeptides can be of any length disclosed within the above ranges and can initiate at any amino acid position represented in the Sequence Listing.

The invention also contemplates "substantially isolated" or "substantially pure" proteins or polypeptides. By a "substantially isolated" or "substantially pure" protein or polypeptide is meant a protein or polypeptide that has been separated from at least some of those components that naturally accompany it. Typically, the protein or polypeptide is substantially isolated or pure when it is at least 60%, by weight, free from the proteins and other naturally-occurring organic molecules with which it is naturally associated *in vivo*. Preferably, the purity of the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight. A substantially isolated or pure protein or polypeptide may be obtained, for example, by extraction from a natural source, by expression of a recombinant nucleic acid encoding the protein or polypeptide, or by chemically synthesizing the protein or polypeptide.

Purity can be measured by any appropriate method, e.g., column chromatography such as immunoaffinity chromatography using an antibody specific for the protein or polypeptide, polyacrylamide gel electrophoresis, or HPLC analysis. A protein or polypeptide is substantially free of naturally associated components when it is separated from at least some of those contaminants that accompany it in its natural state. Thus, a polypeptide that is chemically synthesized or produced in a cellular system different

from the cell from which it naturally originates will be, by definition, substantially free from its naturally associated components. Accordingly, substantially isolated or pure proteins or polypeptides include
5 eukaryotic proteins synthesized in *E. coli*, other prokaryotes, or any other organism in which they do not naturally occur.

5.3 ANTIBODIES TO NHP PRODUCTS

10 Antibodies that specifically recognize one or more epitopes of a NHP, epitopes of conserved variants of a NHP, or peptide fragments of a NHP, are also encompassed by the invention. Such antibodies include, but are not limited to, polyclonal antibodies, monoclonal antibodies
15 (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

20 The antibodies of the invention may be used, for example, in the detection of a NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of a NHP. Such
25 antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP expression product. Additionally, such antibodies can be used in conjunction with gene
30 therapy to, for example, evaluate normal and/or engineered NHP-expressing cells prior to their introduction into a patient. Such antibodies may additionally be used in methods for the inhibition of abnormal NHP activity. Thus, such antibodies may be
35 utilized as a part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with a NHP, a NHP peptide (e.g., one corresponding to a functional domain of a NHP), a truncated NHP polypeptide (a NHP in which one or more domains have been deleted), functional equivalents of a NHP, or mutated variants of a NHP. Such host animals may include, but are not limited to, pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including, but not limited to, Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, chitosan, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and/or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin, or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp.

77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, and IgD, and any subclass thereof. The hybridomas producing the mAbs of this invention may be cultivated *in vitro* or *in vivo*.

- 5 Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger
10 *et al.*, 1984, *Nature*, 312:604-608; Takeda *et al.*, 1985, *Nature*, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric
15 antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patent Nos. 6,114,598, 6,075,181
20 and 5,877,397 and their respective disclosures, which are herein incorporated by reference in their entirety. Also encompassed by the present invention is the use of fully humanized monoclonal antibodies, as described in U.S. Patent No. 6,150,584 and respective disclosures,
25 which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, 1988, *Science* 242:423-426; Huston *et al.*, 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and
30 Ward *et al.*, 1989, *Nature* 341:544-546) can be adapted to produce single chain antibodies against NHP expression products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via
35 an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: F(ab')₂ fragments, which can be produced by pepsin digestion of an antibody molecule; and Fab fragments, which can be generated by reducing the disulfide bridges of F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well-known to those skilled in the art (see, e.g., Greenspan and Bona, 1993, FASEB J. 7:437-444; and Nissinoff, 1991, J. Immunol. 147:2429-2438). For example, antibodies that bind to a NHP domain and competitively inhibit the binding of a NHP to its cognate receptor can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic antibodies, or Fab fragments of such anti-idiotypes, can be used in therapeutic regimens involving a NHP-mediated pathway.

Additionally given the high degree of relatedness of mammalian NHPs, NHP knock-out mice (having never seen a NHP, and thus never been tolerized to a NHP) have a unique utility, as they can be advantageously applied to the generation of antibodies against the disclosed mammalian NHPs (i.e., a NHP will be immunogenic in NHP knock-out animals).

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in

addition to those shown and described herein, will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All
5 cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule
comprising the nucleotide sequence of SEQ ID NO:7.

5

2. The isolated nucleic acid molecule of
claim 1, wherein the molecule comprises the nucleotide
sequence of SEQ ID NO:5.

10

3. The isolated nucleic acid molecule of
claim 2, wherein the molecule comprises the nucleotide
sequence of SEQ ID NO:3.

15

4. The isolated nucleic acid molecule of
claim 3, wherein the molecule comprises the nucleotide
sequence of SEQ ID NO:1.

20

5. An isolated nucleic acid molecule
comprising a nucleotide sequence that:

(a) encodes the amino acid sequence shown in
SEQ ID NO:4; and

(b) hybridizes under highly stringent
conditions to the nucleotide sequence of
SEQ ID NO:3 or the complement thereof.

25

6. An isolated nucleic acid molecule
encoding the amino acid sequence described in SEQ ID
NO:8.

30

7. The isolated nucleic acid molecule of
claim 6, wherein the molecule encodes the amino acid
sequence of SEQ ID NO:6.

35

8. The isolated nucleic acid molecule of
claim 7, wherein the molecule encodes the amino acid
sequence of SEQ ID NO:4.

9. The isolated nucleic acid molecule of claim 8, wherein the molecule encodes the amino acid sequence of SEQ ID NO:2.

5 10. A substantially isolated protein having the activity of the protein shown in SEQ ID NOS:2, 4, 6, or 8, which is encoded by a nucleotide sequence that hybridizes to SEQ ID NO:1, 3, 5, or 7 under highly stringent conditions.

10

15

20

SEQUENCE LISTING

<110> LEXICON GENETICS INCORPORATED

<120> Novel Human Hydroxylases and Polynucleotides Encoding the Same

<130> LEX-0352-PCT

<150> US 60/294,076

<151> 2001-05-29

<160> 8

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1473

<212> DNA

<213> homo sapiens

<400> 1

```

atgcagccag caatgatgat gttttccagt aaatactggg cacggagagg gttttccctg 60
gattcagcag tgcccgaaga gcatcagcta cttggcagct caacactaaa taaacctaac 120
tctggcaaaa atgacgacaa aggcaacaag ggaagcagca aacgtgaagc tgctaccgaa 180
agtggcaaga cagcagttgt tttctccttg aagaatgaag ttggtggatt ggtaaaagca 240
ctgaggctct ttcaggaaaa acgtgtcaac atggttcata ttgaatccag gaaatctcgg 300
cgaagaagtt ctgaggttga aatctttgtg gactgtgagt gtgggaaaaac agaattcaat 360
gagctcattc agttgctgaa atttcaaacc actattgtga cgctgaatcc tccagagAAC 420
atttggacag aggaagaaga gctagaggat gtgccctggg tccctcggaa gatctctgag 480
ttagacaaat gctctcacag agttctcatg tatggttctg agcttgatgc tgaccaccca 540
ggatttaagg acaatgtcta tcgacagaga agaaagtatt ttgtggatgt ggccatgggt 600
tataaatatg gtcagcccat tcccagggtg gagtatactg aagaagaaac taaaacttgg 660
ggtgtgtgat tccgggagct ctocaaactc tatccactc atgcttgccg agagtatttg 720
aaaaacttcc ctctgctgac taaatactgt ggctacagag aggacaatgt gcctcaactc 780
gaagatgtct ccatgtttct gaaagaaagg tctggcttca cggtgaggcc ggtggctgga 840
tacctgagcc cagcagactt tctggcagga ctggcctaca gagtgttcca ctgtacccag 900
tacatccggc atggctcaga tccccctctac accccagaac cagacacatg ccatgaactc 960
ttgggacatg ttccactact tgccgatact aagtttgctc agttttcaca agaaataggt 1020
ctggcgctctc tgggagcatt agatgaagat gttcagaaac tagccacgtg ctatttcttc 1080
acaatcgagt ttggcctttg caagcaagaa gggcaactgc gggcatatgg agcaggactc 1140
ctttcctcca ttggagaatt aaagcacgcc ctttctgaca aggcattgtg gaaagccttt 1200
gacccaaaga caacttgctt acaggaatgc cttatcacca ccttccagga agcctacttt 1260
gtttcagaaa gttttgaaga agccaaagaa aagatgaggg actttgcaaa gtcaattacc 1320
cgtcccttct cagtatactt caatccctac acacagagta ttgaaattct gaaagacacc 1380
agaagtattg aaaatgtggg gcaggacctt cgcagcgact tgaatacagt gtgtgatgct 1440
ttaaacaata tgaaccaata tctggggatt tga
1473

```

<210> 2

<211> 490

<212> PRT

<213> homo sapiens

<400> 2

```

Met Gln Pro Ala Met Met Met Phe Ser Ser Lys Tyr Trp Ala Arg Arg
1      5      10      15
Gly Phe Ser Leu Asp Ser Ala Val Pro Glu Glu His Gln Leu Leu Gly
20      25      30
Ser Ser Thr Leu Asn Lys Pro Asn Ser Gly Lys Asn Asp Asp Lys Gly
35      40      45
Asn Lys Gly Ser Ser Lys Arg Glu Ala Ala Thr Glu Ser Gly Lys Thr
50      55      60
Ala Val Val Phe Ser Leu Lys Asn Glu Val Gly Gly Leu Val Lys Ala

```

65					70					75				80	
Leu	Arg	Leu	Phe	Gln	Glu	Lys	Arg	Val	Asn	Met	Val	His	Ile	Glu	Ser
				85					90					95	
Arg	Lys	Ser	Arg	Arg	Arg	Ser	Ser	Glu	Val	Glu	Ile	Phe	Val	Asp	Cys
			100					105					110		
Glu	Cys	Gly	Lys	Thr	Glu	Phe	Asn	Glu	Leu	Ile	Gln	Leu	Leu	Lys	Phe
		115					120					125			
Gln	Thr	Thr	Ile	Val	Thr	Leu	Asn	Pro	Pro	Glu	Asn	Ile	Trp	Thr	Glu
		130					135				140				
Glu	Glu	Glu	Leu	Glu	Asp	Val	Pro	Trp	Phe	Pro	Arg	Lys	Ile	Ser	Glu
				150						155					160
Leu	Asp	Lys	Cys	Ser	His	Arg	Val	Leu	Met	Tyr	Gly	Ser	Glu	Leu	Asp
				165					170					175	
Ala	Asp	His	Pro	Gly	Phe	Lys	Asp	Asn	Val	Tyr	Arg	Gln	Arg	Arg	Lys
			180					185					190		
Tyr	Phe	Val	Asp	Val	Ala	Met	Gly	Tyr	Lys	Tyr	Gly	Gln	Pro	Ile	Pro
		195					200					205			
Arg	Val	Glu	Tyr	Thr	Glu	Glu	Glu	Thr	Lys	Thr	Trp	Gly	Val	Val	Phe
		210					215				220				
Arg	Glu	Leu	Ser	Lys	Leu	Tyr	Pro	Thr	His	Ala	Cys	Arg	Glu	Tyr	Leu
		225			230					235					240
Lys	Asn	Phe	Pro	Leu	Leu	Thr	Lys	Tyr	Cys	Gly	Tyr	Arg	Glu	Asp	Asn
				245					250					255	
Val	Pro	Gln	Leu	Glu	Asp	Val	Ser	Met	Phe	Leu	Lys	Glu	Arg	Ser	Gly
			260					265					270		
Phe	Thr	Val	Arg	Pro	Val	Ala	Gly	Tyr	Leu	Ser	Pro	Arg	Asp	Phe	Leu
		275					280					285			
Ala	Gly	Leu	Ala	Tyr	Arg	Val	Phe	His	Cys	Thr	Gln	Tyr	Ile	Arg	His
		290				295					300				
Gly	Ser	Asp	Pro	Leu	Tyr	Thr	Pro	Glu	Pro	Asp	Thr	Cys	His	Glu	Leu
				310						315					320
Leu	Gly	His	Val	Pro	Leu	Leu	Ala	Asp	Pro	Lys	Phe	Ala	Gln	Phe	Ser
				325					330					335	
Gln	Glu	Ile	Gly	Leu	Ala	Ser	Leu	Gly	Ala	Ser	Asp	Glu	Asp	Val	Gln
			340					345					350		
Lys	Leu	Ala	Thr	Cys	Tyr	Phe	Phe	Thr	Ile	Glu	Phe	Gly	Leu	Cys	Lys
		355					360					365			
Gln	Glu	Gly	Gln	Leu	Arg	Ala	Tyr	Gly	Ala	Gly	Leu	Leu	Ser	Ser	Ile
		370				375					380				
Gly	Glu	Leu	Lys	His	Ala	Leu	Ser	Asp	Lys	Ala	Cys	Val	Lys	Ala	Phe
				390						395					400
Asp	Pro	Lys	Thr	Thr	Cys	Leu	Gln	Glu	Cys	Leu	Ile	Thr	Thr	Phe	Gln
				405					410					415	
Glu	Ala	Tyr	Phe	Val	Ser	Glu	Ser	Phe	Glu	Glu	Ala	Lys	Glu	Lys	Met
			420					425					430		
Arg	Asp	Phe	Ala	Lys	Ser	Ile	Thr	Arg	Pro	Phe	Ser	Val	Tyr	Phe	Asn
		435					440					445			
Pro	Tyr	Thr	Gln	Ser	Ile	Glu	Ile	Leu	Lys	Asp	Thr	Arg	Ser	Ile	Glu
		450				455					460				
Asn	Val	Val	Gln	Asp	Leu	Arg	Ser	Asp	Leu	Asn	Thr	Val	Cys	Asp	Ala
				470						475					480
Leu	Asn	Lys	Met	Asn	Gln	Tyr	Leu	Gly	Ile						
				485					490						

<210> 3

<211> 1461

<212> DNA

<213> homo sapiens

<400> 3

atgatgatgt tttccagtaa atactgggca cggagagggt tttccctgga ttcagcagtg 60
 cccgaagagc atcagctact tggcagctca acactaaata aacctaactc tggcaaaaat 120


```

gacgacaaag gcaacaaggg aagcagcaaa cgtgaagctg ctaccgaaag tggcaagaca 180
gcagttgttt tctccttgaa gaatgaagtt ggtggattgg taaaagcact gaggtctttt 240
caggaaaaac gtgtcaacat ggttcatatt gaatccagga aatctcggcg aagaagttct 300
gaggttgaaa tcttttgga ctgtgagtg gggaaaacag aattcaatga gctcattcag 360
ttgctgaaat ttcaaaccac tattgtgacg ctgaatcctc cagagaacat ttggacagag 420
gaagaagagc tagaggatgt gccctgggtc cctcggaaga tctctgagtt agacaaatgc 480
tctcacagag ttctcatgta tggttctgag cttgatgctg accaccagg atttaaggac 540
aatgtctatc gacagagaag aaagtatttt gtggatgtgg ccatgggtta taaatatggt 600
cagcccatc ccagggtgga gtatactgaa gaagaaacta aaacttgggg tgtgtattc 660
cgggagctct ccaaactcta tcccactcat gcttgccgag agtatttgaa aaacttccct 720
ctgctgacta aatactgtgg ctacagagag gacaatgtgc ctcaactcga agatgtctcc 780
atgtttctga aagaaagggtc tggcttcacg gtgaggccgg tggctggata cctgagccca 840
cgagactttc tggcaggact ggcctacaga gtgttccact gtaccagta catccggcat 900
ggctcagatc cctctacac cccagaacca gacacatgcc atgaactctt gggacatgtt 960
ccactacttg cggatcctaa gtttgctcag tttcacaaag aaataggtct ggcgtctctg 1020
ggagcatcag atgaagatgt tcagaaacta gccacgtgct atttcttcac aatcgagttt 1080
ggcctttgca agcaagaagg gcaactgagg gcatatggag caggactcct ttcctccatt 1140
ggagaattaa agcacgccct ttctgacaag gcatgtgtga aagcctttga cccaaagaca 1200
acttgcttac aggaatgcct taccaccacc ttccagggaag cctactttgt ttcagaaagt 1260
tttgaagaag ccaagaagaaa gatgagggac tttgcaaaagt caattaccg tcccttctca 1320
gtatacttca atccctacac acagagtatt gaaattctga aagacaccag aagtattgaa 1380
aatgtggtgc aggaccttcg cagcgacttg aatacagtgt gtgatgcttt aaacaaaatg 1440
aaccaatata tggggatttg a                                     1461

```

<210> 4

<211> 486

<212> PRT

<213> homo sapiens

<400> 4

```

Met Met Met Phe Ser Ser Lys Tyr Trp Ala Arg Arg Gly Phe Ser Leu
 1          5          10          15
Asp Ser Ala Val Pro Glu Glu His Gln Leu Leu Gly Ser Ser Thr Leu
          20          25          30
Asn Lys Pro Asn Ser Gly Lys Asn Asp Asp Lys Gly Asn Lys Gly Ser
          35          40          45
Ser Lys Arg Glu Ala Ala Thr Glu Ser Gly Lys Thr Ala Val Val Phe
          50          55          60
Ser Leu Lys Asn Glu Val Gly Gly Leu Val Lys Ala Leu Arg Leu Phe
          65          70          75          80
Gln Glu Lys Arg Val Asn Met Val His Ile Glu Ser Arg Lys Ser Arg
          85          90          95
Arg Arg Ser Ser Glu Val Glu Ile Phe Val Asp Cys Glu Cys Gly Lys
          100          105          110
Thr Glu Phe Asn Glu Leu Ile Gln Leu Leu Lys Phe Gln Thr Thr Ile
          115          120          125
Val Thr Leu Asn Pro Pro Glu Asn Ile Trp Thr Glu Glu Glu Glu Leu
          130          135          140
Glu Asp Val Pro Trp Phe Pro Arg Lys Ile Ser Glu Leu Asp Lys Cys
          145          150          155          160
Ser His Arg Val Leu Met Tyr Gly Ser Glu Leu Asp Ala Asp His Pro
          165          170          175
Gly Phe Lys Asp Asn Val Tyr Arg Gln Arg Arg Lys Tyr Phe Val Asp
          180          185          190
Val Ala Met Gly Tyr Lys Tyr Gly Gln Pro Ile Pro Arg Val Glu Tyr
          195          200          205
Thr Glu Glu Glu Thr Lys Thr Trp Gly Val Val Phe Arg Glu Leu Ser
          210          215          220
Lys Leu Tyr Pro Thr His Ala Cys Arg Glu Tyr Leu Lys Asn Phe Pro
          225          230          235          240
Leu Leu Thr Lys Tyr Cys Gly Tyr Arg Glu Asp Asn Val Pro Gln Leu
          245          250          255
Glu Asp Val Ser Met Phe Leu Lys Glu Arg Ser Gly Phe Thr Val Arg

```

Pro	Val	Ala	260	Gly	Tyr	Leu	Ser	Pro	265	Arg	Asp	Phe	Leu	Ala	270	Gly	Leu	Ala
		275						280						285				
Tyr	Arg	Val	Phe	His	Cys	Thr	Gln	Tyr	Ile	Arg	His	Gly	Ser	Asp	Pro			
	290					295					300							
Leu	Tyr	Thr	Pro	Glu	Pro	Asp	Thr	Cys	His	Glu	Leu	Leu	Gly	His	Val			
305					310					315					320			
Pro	Leu	Leu	Ala	Asp	Pro	Lys	Phe	Ala	Gln	Phe	Ser	Gln	Glu	Ile	Gly			
			325					330						335				
Leu	Ala	Ser	Leu	Gly	Ala	Ser	Asp	Glu	Asp	Val	Gln	Lys	Leu	Ala	Thr			
		340						345					350					
Cys	Tyr	Phe	Phe	Thr	Ile	Glu	Phe	Gly	Leu	Cys	Lys	Gln	Glu	Gly	Gln			
	355					360					365							
Leu	Arg	Ala	Tyr	Gly	Ala	Gly	Leu	Leu	Ser	Ser	Ile	Gly	Glu	Leu	Lys			
	370					375					380							
His	Ala	Leu	Ser	Asp	Lys	Ala	Cys	Val	Lys	Ala	Phe	Asp	Pro	Lys	Thr			
385					390					395					400			
Thr	Cys	Leu	Gln	Glu	Cys	Leu	Ile	Thr	Thr	Phe	Gln	Glu	Ala	Tyr	Phe			
			405					410						415				
Val	Ser	Glu	Ser	Phe	Glu	Glu	Ala	Lys	Glu	Lys	Met	Arg	Asp	Phe	Ala			
		420						425					430					
Lys	Ser	Ile	Thr	Arg	Pro	Phe	Ser	Val	Tyr	Phe	Asn	Pro	Tyr	Thr	Gln			
	435					440					445							
Ser	Ile	Glu	Ile	Leu	Lys	Asp	Thr	Arg	Ser	Ile	Glu	Asn	Val	Val	Gln			
	450					455					460							
Asp	Leu	Arg	Ser	Asp	Leu	Asn	Thr	Val	Cys	Asp	Ala	Leu	Asn	Lys	Met			
465					470			475							480			
Asn	Gln	Tyr	Leu	Gly	Ile													
			485															

<210> 5

<211> 1458

<212> DNA

<213> homo sapiens

<400> 5

atgatgtttt	ccagtaaata	ctgggcacgg	agaggggtttt	ccctggattc	agcagtgcc	60
gaagagcatc	agctacttgg	cagctcaaca	ctaaataaac	ctaactctgg	caaaaatgac	120
gacaaaggca	acaagggaag	cagcaaacgt	gaagctgcta	ccgaaagtgg	caagacagca	180
gttggtttct	ccttgaagaa	tgaagttggt	ggattggtaa	aagcactgag	gctctttcag	240
gaaaaacgtg	tcaacatggt	tcatattgaa	tccaggaaat	ctcggcgaag	aagttctgag	300
gttgaaatct	ttgtggactg	tgagtgtggg	aaaacagaat	tcaatgagct	cattcagttg	360
ctgaaatttc	aaaccactat	tgtgacgctg	aatcctccag	agaacatttg	gacagaggaa	420
gaagagctag	aggatgtgcc	ctggttccct	cggaagatct	ctgagttaga	caaagtctct	480
cacagagtgc	tcatgtatgg	ttctgagctt	gatgctgacc	acccaggatt	taaggacaat	540
gtctatcgac	agagaagaaa	gtattttgtg	gatgtggcca	tgggttataa	atatggtcag	600
cccattccca	gggtggagta	tactgaagaa	gaaactaaaa	cttgggggtg	tgtattccgg	660
gagctctcca	aactctatcc	cactcatgct	tgccgagagt	atttgaaaaa	cttcctctcg	720
ctgactaaa	actgtggcta	cagagaggac	aatgtgcctc	aactcgaaga	tgtctccatg	780
tttctgaaag	aaaggtctgg	cttcacggtg	aggccggtgg	ctggatacct	gagcccacga	840
gactttcttg	caggactggc	ctacagagtg	ttccactgta	cccagtacat	ccggcatggc	900
tcagatcccc	tctacacccc	agaaccagac	acatgccatg	aactcttggg	acatgttcca	960
ctacttgctg	atcctaagtt	tgctcagttt	tcacaagaaa	taggtctggc	gtctctggga	1020
gcatcagatg	aagatgttca	gaaactagcc	acgtgctatt	tcttcacaat	cgagtttggc	1080
ctttgcaagc	aagaagggca	actgcgggca	tatggagcag	gactcctttc	ctccattgga	1140
gaattaaagc	acgccctttc	tgacaaggca	tgtgtgaaag	cctttgaccc	aaagacaact	1200
tgcttacagg	aatgccttat	caccaccttc	caggaagcct	actttgtttc	agaaagtgtt	1260
gaagaagcca	aagaaaagat	gagggacttt	gcaaagtcaa	ttaccggtcc	cttctcagta	1320
tacttcaatc	cctacacaca	gagtattgaa	attctgaaag	acaccagaag	tattgaaaat	1380
gtgggtgcagg	accttcgcag	cgacttgaat	acagtgtgtg	atgcttttaa	caaaatgaac	1440
caatatctgg	ggatttga					1458

<210> 6
 <211> 485
 <212> PRT
 <213> homo sapiens

<400> 6

Met	Met	Phe	Ser	Ser	Lys	Tyr	Trp	Ala	Arg	Arg	Gly	Phe	Ser	Leu	Asp
1				5					10					15	
Ser	Ala	Val	Pro	Glu	Glu	His	Gln	Leu	Leu	Gly	Ser	Ser	Thr	Leu	Asn
			20					25					30		
Lys	Pro	Asn	Ser	Gly	Lys	Asn	Asp	Asp	Lys	Gly	Asn	Lys	Gly	Ser	Ser
		35					40					45			
Lys	Arg	Glu	Ala	Ala	Thr	Glu	Ser	Gly	Lys	Thr	Ala	Val	Val	Phe	Ser
	50					55				60					
Leu	Lys	Asn	Glu	Val	Gly	Gly	Leu	Val	Lys	Ala	Leu	Arg	Leu	Phe	Gln
65					70					75					80
Glu	Lys	Arg	Val	Asn	Met	Val	His	Ile	Glu	Ser	Arg	Lys	Ser	Arg	Arg
				85					90					95	
Arg	Ser	Ser	Glu	Val	Glu	Ile	Phe	Val	Asp	Cys	Glu	Cys	Gly	Lys	Thr
			100					105					110		
Glu	Phe	Asn	Glu	Leu	Ile	Gln	Leu	Leu	Lys	Phe	Gln	Thr	Thr	Ile	Val
		115				120						125			
Thr	Leu	Asn	Pro	Pro	Glu	Asn	Ile	Trp	Thr	Glu	Glu	Glu	Glu	Leu	Glu
		130				135					140				
Asp	Val	Pro	Trp	Phe	Pro	Arg	Lys	Ile	Ser	Glu	Leu	Asp	Lys	Cys	Ser
145					150					155					160
His	Arg	Val	Leu	Met	Tyr	Gly	Ser	Glu	Leu	Asp	Ala	Asp	His	Pro	Gly
			165						170					175	
Phe	Lys	Asp	Asn	Val	Tyr	Arg	Gln	Arg	Arg	Lys	Tyr	Phe	Val	Asp	Val
			180					185					190		
Ala	Met	Gly	Tyr	Lys	Tyr	Gly	Gln	Pro	Ile	Pro	Arg	Val	Glu	Tyr	Thr
		195					200					205			
Glu	Glu	Glu	Thr	Lys	Thr	Trp	Gly	Val	Val	Phe	Arg	Glu	Leu	Ser	Lys
		210				215					220				
Leu	Tyr	Pro	Thr	His	Ala	Cys	Arg	Glu	Tyr	Leu	Lys	Asn	Phe	Pro	Leu
225					230					235					240
Leu	Thr	Lys	Tyr	Cys	Gly	Tyr	Arg	Glu	Asp	Asn	Val	Pro	Gln	Leu	Glu
			245						250					255	
Asp	Val	Ser	Met	Phe	Leu	Lys	Glu	Arg	Ser	Gly	Phe	Thr	Val	Arg	Pro
			260					265					270		
Val	Ala	Gly	Tyr	Leu	Ser	Pro	Arg	Asp	Phe	Leu	Ala	Gly	Leu	Ala	Tyr
		275					280					285			
Arg	Val	Phe	His	Cys	Thr	Gln	Tyr	Ile	Arg	His	Gly	Ser	Asp	Pro	Leu
		290				295					300				
Tyr	Thr	Pro	Glu	Pro	Asp	Thr	Cys	His	Glu	Leu	Leu	Gly	His	Val	Pro
305					310					315					320
Leu	Leu	Ala	Asp	Pro	Lys	Phe	Ala	Gln	Phe	Ser	Gln	Glu	Ile	Gly	Leu
			325						330					335	
Ala	Ser	Leu	Gly	Ala	Ser	Asp	Glu	Asp	Val	Gln	Lys	Leu	Ala	Thr	Cys
			340					345					350		
Tyr	Phe	Phe	Thr	Ile	Glu	Phe	Gly	Leu	Cys	Lys	Gln	Glu	Gly	Gln	Leu
		355					360					365			
Arg	Ala	Tyr	Gly	Ala	Gly	Leu	Leu	Ser	Ser	Ile	Gly	Glu	Leu	Lys	His
		370				375					380				
Ala	Leu	Ser	Asp	Lys	Ala	Cys	Val	Lys	Ala	Phe	Asp	Pro	Lys	Thr	Thr
385					390					395					400
Cys	Leu	Gln	Glu	Cys	Leu	Ile	Thr	Thr	Phe	Gln	Glu	Ala	Tyr	Phe	Val
			405						410					415	
Ser	Glu	Ser	Phe	Glu	Glu	Ala	Lys	Glu	Lys	Met	Arg	Asp	Phe	Ala	Lys
			420					425					430		
Ser	Ile	Thr	Arg	Pro	Phe	Ser	Val	Tyr	Phe	Asn	Pro	Tyr	Thr	Gln	Ser
		435					440					445			
Ile	Glu	Ile	Leu	Lys	Asp	Thr	Arg	Ser	Ile	Glu	Asn	Val	Val	Gln	Asp

450 455 460
 Leu Arg Ser Asp Leu Asn Thr Val Cys Asp Ala Leu Asn Lys Met Asn
 465 470 475 480
 Gln Tyr Leu Gly Ile
 485

<210> 7
 <211> 1455
 <212> DNA
 <213> homo sapiens

<400> 7
 atgtttttccca gtaaataactg ggcacggaga ggggttttccc tggattcagc agtgcccga 60
 gagcatcagc tacttggcag ctcaacacta aataaaccta actctggcaa aaatgacgac 120
 aaaggcaaca aggggaagcag caaacgtgaa gctgctaccg aaagtggcaa gacagcagtt 180
 gttttctcct tgaagaatga agttgggtgga ttggtaaaaag cactgagggt ctttcaggaa 240
 aaacgtgtca acatgggttca tattgaatcc aggaaatctc ggcaagaag ttctgaggtt 300
 gaaatctttg tggactgtga gtgtgggaaa acagaattca atgagctcat tcagttgctg 360
 aaatttcaaa ccactattgt gacgctgaat cctccagaga acatttggac agaggaagaa 420
 gagctagagg atgtgccctg gttccctcgg aagatctctg agttagacaa atgctctcac 480
 agagtcttcca tgtatggttc tgagcttgat gctgaccacc caggatttaa ggacaatgtc 540
 tatcgacaga gaagaaagta ttttgtggat gtggccatgg gttataaata tggtcagccc 600
 attcccaggg tggagtatac tgaagaagaa actaaaactt ggggtgttgt attccgggag 660
 ctctccaaac tctatccccc tcatgcttgc cgagagtatt tgaaaaactt cctctgctg 720
 actaaatact gtggctacag agaggacaat gtgcctcaac tcgaagatgt ctccatgttt 780
 ctgaaagaaa ggtctggctt cacggtgag cgggtggctg gatacctgag ccacagagac 840
 tttctggcag gactggccta cagagtgttc cactgtacc cagacatccg gcatggctca 900
 gatccccctc acaccccgaga accagacaca tgccatgaac tcttgggaca tgttccacta 960
 cttgcggtatc ctaagtttgc tcagttttca caagaaatag gtctggcgctc tctgggagca 1020
 tcagatgaag atgttcagaa actagccacg tgctatttct tcacaatcga gtttggcctt 1080
 tgcaagcaag aagggcaact gcgggcatat ggagcaggac tcctttcctc cattggagaa 1140
 ttaaagcacg ccctttctga caaggcatgt gtgaaagcct ttgacccaaa gacaacttgc 1200
 ttacaggaat gccttatcac caccttcag gaagcctact ttgtttcaga aagttttgaa 1260
 gaagccaaag aaaagatgag ggactttgca aagtcaatta cccgtccctt ctcaagtatac 1320
 ttcaatccct acacacagag tattgaaatt ctgaaagaca ccagaagtat tgaaaatgtg 1380
 gtgcaggacc ttcgcagcga cttgaatata gtgtgtgatg ctttaaacaa aatgaaccaa 1440
 tatctgggga tttga 1455

<210> 8
 <211> 484
 <212> PRT
 <213> homo sapiens

<400> 8
 Met Phe Ser Ser Lys Tyr Trp Ala Arg Arg Gly Phe Ser Leu Asp Ser
 1 5 10 15
 Ala Val Pro Glu Glu His Gln Leu Leu Gly Ser Ser Thr Leu Asn Lys
 20 25 30
 Pro Asn Ser Gly Lys Asn Asp Asp Lys Gly Asn Lys Gly Ser Ser Lys
 35 40 45
 Arg Glu Ala Ala Thr Glu Ser Gly Lys Thr Ala Val Val Phe Ser Leu
 50 55 60
 Lys Asn Glu Val Gly Gly Leu Val Lys Ala Leu Arg Leu Phe Gln Glu
 65 70 75 80
 Lys Arg Val Asn Met Val His Ile Glu Ser Arg Lys Ser Arg Arg
 85 90 95
 Ser Ser Glu Val Glu Ile Phe Val Asp Cys Glu Cys Gly Lys Thr Glu
 100 105 110
 Phe Asn Glu Leu Ile Gln Leu Leu Lys Phe Gln Thr Thr Ile Val Thr
 115 120 125
 Leu Asn Pro Pro Glu Asn Ile Trp Thr Glu Glu Glu Glu Leu Glu Asp
 130 135 140

Val	Pro	Trp	Phe	Pro	Arg	Lys	Ile	Ser	Glu	Leu	Asp	Lys	Cys	Ser	His
145					150					155					160
Arg	Val	Leu	Met	Tyr	Gly	Ser	Glu	Leu	Asp	Ala	Asp	His	Pro	Gly	Phe
			165						170					175	
Lys	Asp	Asn	Val	Tyr	Arg	Gln	Arg	Arg	Lys	Tyr	Phe	Val	Asp	Val	Ala
		180						185					190		
Met	Gly	Tyr	Lys	Tyr	Gly	Gln	Pro	Ile	Pro	Arg	Val	Glu	Tyr	Thr	Glu
	195						200					205			
Glu	Glu	Thr	Lys	Thr	Trp	Gly	Val	Val	Phe	Arg	Glu	Leu	Ser	Lys	Leu
	210					215					220				
Tyr	Pro	Thr	His	Ala	Cys	Arg	Glu	Tyr	Leu	Lys	Asn	Phe	Pro	Leu	Leu
225					230					235					240
Thr	Lys	Tyr	Cys	Gly	Tyr	Arg	Glu	Asp	Asn	Val	Pro	Gln	Leu	Glu	Asp
			245						250					255	
Val	Ser	Met	Phe	Leu	Lys	Glu	Arg	Ser	Gly	Phe	Thr	Val	Arg	Pro	Val
		260						265					270		
Ala	Gly	Tyr	Leu	Ser	Pro	Arg	Asp	Phe	Leu	Ala	Gly	Leu	Ala	Tyr	Arg
	275						280					285			
Val	Phe	His	Cys	Thr	Gln	Tyr	Ile	Arg	His	Gly	Ser	Asp	Pro	Leu	Tyr
	290					295					300				
Thr	Pro	Glu	Pro	Asp	Thr	Cys	His	Glu	Leu	Leu	Gly	His	Val	Pro	Leu
305					310					315					320
Leu	Ala	Asp	Pro	Lys	Phe	Ala	Gln	Phe	Ser	Gln	Glu	Ile	Gly	Leu	Ala
			325						330					335	
Ser	Leu	Gly	Ala	Ser	Asp	Glu	Asp	Val	Gln	Lys	Leu	Ala	Thr	Cys	Tyr
		340					345						350		
Phe	Phe	Thr	Ile	Glu	Phe	Gly	Leu	Cys	Lys	Gln	Glu	Gly	Gln	Leu	Arg
	355					360						365			
Ala	Tyr	Gly	Ala	Gly	Leu	Leu	Ser	Ser	Ile	Gly	Glu	Leu	Lys	His	Ala
	370					375				380					
Leu	Ser	Asp	Lys	Ala	Cys	Val	Lys	Ala	Phe	Asp	Pro	Lys	Thr	Thr	Cys
385					390					395					400
Leu	Gln	Glu	Cys	Leu	Ile	Thr	Thr	Phe	Gln	Glu	Ala	Tyr	Phe	Val	Ser
			405						410					415	
Glu	Ser	Phe	Glu	Glu	Ala	Lys	Glu	Lys	Met	Arg	Asp	Phe	Ala	Lys	Ser
		420						425					430		
Ile	Thr	Arg	Pro	Phe	Ser	Val	Tyr	Phe	Asn	Pro	Tyr	Thr	Gln	Ser	Ile
	435						440					445			
Glu	Ile	Leu	Lys	Asp	Thr	Arg	Ser	Ile	Glu	Asn	Val	Val	Gln	Asp	Leu
	450					455				460					
Arg	Ser	Asp	Leu	Asn	Thr	Val	Cys	Asp	Ala	Leu	Asn	Lys	Met	Asn	Gln
465					470					475					480
Tyr	Leu	Gly	Ile												

THIS PAGE BLANK (USPTO)